arXiv:1209.2358v1 [math.QA] 11 Sep 2012

SOME CATEGORICAL REPRESENTATIONS OF THE MODULAR GROUP

BENJAMIN COOPER

Abstract. Examples of $\mathrm{SL}(2,\mathbb{Z})$ actions on differential graded categories are defined and explored.

1. Introduction

The study of topological field theories leads to an interest in group actions on categories. In some formulations, the codimension 2 portion of an n-dimensional topological field theory associates to each oriented $(n-2)$-manifold Σ a differential graded category $\mathcal{D}(\Sigma)$,

$$\Sigma \rightsquigarrow \mathcal{D}(\Sigma),$$

upon which the mapping class group of the manifold $\Gamma(\Sigma)$ acts by functors $F_g : \mathcal{D}(\Sigma) \to \mathcal{D}(\Sigma)$ for $g \in \Gamma(\Sigma)$. For discussion see [12, 17].

It is believed that there are many examples of such topological field theories when $n = 4$. In particular, there should be many examples of dg categories supporting actions of mapping class groups of surfaces. Unfortunately, while there are a number of braid group representations of this type [2, 13, 19, 18], there are few examples of such mapping class group representations for surfaces of genus greater than zero (for an exception, see [14]). Ideally, one would like a body of examples to draw upon when investigating a new phenomenon.

This paper presents a novel means by which some categories associated to the Khovanov homology of knots and links can be used to produce differential graded categories which are representations of the modular group $\mathrm{SL}(2,\mathbb{Z})$; the mapping class group of the torus.

In section 2, definitions related to knot homology, group actions and differential graded categories are recalled. In section 3, we review necessary information about braid groups and mapping class groups. In section 4, we produce a countable family of categories $\mathcal{R}_{n,k}$ supporting $\mathrm{SL}(2,\mathbb{Z})$ actions. Section 5 explores the simplest example.

2. Algebraic Background

In this section we collect some background information which is necessary to understand the remainder of the paper. We begin by defining the differential graded categories $\mathrm{Kom}^*(n)$ and recall the existence of certain idempotent chain complexes $P_{n,k}$ within $\mathrm{Kom}^*(n)$. The definition of a group action on a dg category and the definition of the orbit construction in the theory of dg categories are also given.

2.1. **Cobordisms.** Dror Bar-Natan's graphical formulation [4] of the Khovanov categorification [11, 12] consists of a series of categories $\mathrm{Cob}(n)$ and $\mathrm{Kom}(n)$. The construction in section 4 will take place within categories related to $\mathrm{Kom}(n)$. These categories are essentially the same as those appearing in [6, 7] with the exception that we adopt the grading convention appearing in Rozansky [16].

There is a pre-additive category $\text{Pre-Cob}(n)$ whose objects are isotopy classes of formally $\mathbb{Z}$-graded 1-manifolds with $2n$ boundary points. There is a functor

$$q : \text{Pre-Cob}(n) \to \text{Pre-Cob}(n)$$

which increases the grading of all objects by one. The morphisms of the category $\text{Pre-Cob}(n)$ are given by linear combinations of isotopy classes of orientable cobordisms. The internal *degree* of a cobordism $C : q^i A \to q^j B$ is given by

$$\deg(C) = \deg_\chi(C) + \deg_q(C)$$

where the topological degree $\deg_\chi(C) = \chi(C) - n$ is given by the Euler characteristic of C and the q-degree $\deg_q(C) = j - i$ is given by the relative difference in q-gradings.

We impose the relations implied by the requirement that

$$\bigcirc \quad \cong \quad q\emptyset \oplus q^{-1}\emptyset$$

using maps of internal degree zero (see [3]). In doing so, we obtain a new category $\mathrm{Cob}(n)$ as a quotient of the additive closure $\mathrm{Mat}(\text{Pre-Cob}(n))$ of $\text{Pre-Cob}(n)$. There is a composition

$$\otimes : \mathrm{Cob}(n) \times \mathrm{Cob}(n) \to \mathrm{Cob}(n)$$

obtained by gluing of cobordisms.

Denote by $\text{Pre-Kom}(n)$ the category $\mathrm{Cob}(n)$ with an additional $\frac{1}{2}\mathbb{Z}$-grading t and let $\mathrm{Kom}(n)$ be the category of convolutions in $\text{Pre-Kom}(n)$. The objects of $\mathrm{Kom}(n)$ are chain complexes

$$(\oplus_i D_i, d_{ij})$$

with $D_i \in \text{Pre-Kom}(n)$ and $d_{ij} : D_i \to D_j$. The matrix $d = (d_{ij})$ must be lower triangular and square to zero. Each complex is required to be bounded below in sufficiently negative degree and possess finitely many terms in any given $(t^{\frac{1}{2}}, q)$-degree.

Associated to a crossing is the complex,



This definition extends to yield a chain complex in $\mathrm{Kom}(n)$ for each framed (n, n)-tangle. Any two chain complexes associated to tangles which differ by Reidemeister moves 2 or 3 are homotopy equivalent. The first Reidemeister move results in a degree shift, see property 2 below.

The Grothendieck group $\mathrm{K}_0(\mathrm{Kom}(n))$ is defined by the requirement that $\mathrm{K}_0(q) = -q$ and $\mathrm{K}_0(t^{\frac{1}{2}}) = q^{\frac{1}{2}}$. When this is done, the skein relation above becomes the Kauffman bracket.

Each category $\mathrm{Kom}(n)$ can be extended to form a differential graded category $\mathrm{Kom}^*(n)$ which has the same objects as $\mathrm{Kom}(n)$, but morphisms are given by allowing maps of all homological degrees. If $f \in \mathrm{Hom}^m(A, B)$ is a map of homological degree m then $d(f) = [d, f] \in \mathrm{Hom}^{m+1}(A, B)$. Thus, each collection $\mathrm{Hom}^*(A, B)$ of morphisms is a chain complex and the differential d is a derivation with respect to composition of maps.

2.2. **Projectors.** In [6], V. Krushkal and the author defined a special idempotent element $P_n \in \mathrm{Kom}(n)$ which categorified the Jones-Wenzl projector. See also [9, 15].

Theorem 2.3. *There exists a chain complex $P_n \in \mathrm{Kom}(n)$ called the* universal projector *which satisfies*

1. *P_n is positively graded with differential having internal degree zero.*
2. *The identity diagram appears only in homological degree zero and only once.*
3. *The chain complex P_n is contractible under turnbacks.*

These three properties characterize P_n uniquely up to homotopy.

In a more recent paper [7], M. Hogancamp and the author constructed a series of objects $P_{n,k} \in \mathrm{Kom}(n)$ where $k = n, n-2, n-4, \ldots, n \,(\mathrm{mod}\, 2)$. These higher order projectors are characterized uniquely up to homotopy by properties analogous to those mentioned in the theorem above, see section 10 of [7].

If the projector $P_{n,k}$ is represented by a box then it satisfies the two properties below.

Property 1. *(drags through tangles)*



Property 2. *(absorbs twists)*

 $\simeq \quad t^{\frac{1}{2}k^2}q^k$  and  $\simeq \quad t^{-\frac{1}{2}k^2}q^{-k}$ 

Note that it is necessary for the skein relation to be the one defined in section 2.1 for these degree shifts to accurate. When $n = 1$ and $k = 1$ the second property determines the grading shift associated to the first Reidemeister move.

2.4. **Group Actions On Categories.** If G is a group and $\mathcal{C}$ is a differential graded category then an action of G on $\mathcal{C}$ is a homotopy homomorphism from the group G to the monoid of dg functors from $\mathcal{C}$ to $\mathcal{C}$.

$$\varphi : G \to \mathrm{End}(\mathcal{C}) \quad \text{and} \quad \varphi(a \cdot b) \simeq \varphi(a) \circ \varphi(b)$$

If $G = \langle S : R \rangle$ is a presentation for G then a group action is obtained by specifying functors $F_s, F_{s^{-1}} : \mathcal{C} \to \mathcal{C}$ for each generator $s \in S$, and homotopy equivalences

$$F_{s_1} \circ F_{s_2} \circ \cdots \circ F_{s_N} \simeq F_{s'_1} \circ F_{s'_2} \circ \cdots \circ F_{s'_M}$$

whenever the words $s_1 s_2 \cdots s_N$ and $s'_1 s'_2 \cdots s'_M$ coincide by virtue of the relations in R.

Such actions are sometimes called *weak*. An action is strong when the equivalences associated to applications of the relations are uniquely determined.

In a different language, a stratification of EG is a presentation for an ∞-category in which all morphisms are equivalences. A weak action on $\mathcal{C}$ is a functor defined on the 2-skeleton: $EG^{(2)} \to \mathrm{End}(\mathcal{C})$. An action is strong if it satisfies coherence relations which correspond to attaching maps in a choice of basis for $\pi_3(EG^{(2)})$.

We will only prove that the actions defined here are weak.

2.5. **Orbit Categories.** Suppose that $\mathcal{C}$ is a differential graded category and $F : \mathcal{C} \to \mathcal{C}$ is an auto-equivalence. The dg orbit category $\mathcal{C}/F$ has the same objects as the category $\mathcal{C}$. Given any two objects X and Y in $\mathcal{C}$ the morphisms from X to Y are given by

$$\mathrm{Hom}_{\mathcal{C}/F}(X, Y) = \mathrm{colim}_p \bigoplus_{n \geq 0} \mathrm{Hom}_{\mathcal{C}}(F^n X, F^p Y),$$

see [10]. One composes two maps $f : Y \to F^q Z$ and $g : X \to F^p Y$ by first shifting f by the appropriate power of F, $(F^p f) \circ g$.

There is a functor $\pi : \mathcal{C} \to \mathcal{C}/F$ which is universal with respect to the equivalence,

$$\pi \circ F \simeq \pi.$$

We will denote objects of $\mathcal{C}/F$ by πX where $X \in \mathcal{C}$; the functor F being understood from context.

Example. If $\mathcal{C}$ is a dg subcategory of $\mathrm{Kom}^*(n)$ which is closed under degree shifts then each object has both a t-grading and a q-grading and there is a functor

$$F = t^N q^M : \mathcal{C} \to \mathcal{C}$$

which shifts the t-grading and q-grading by N and M respectively. The relation above implies that F acts by identity on the orbit category,

$$\pi F(X) \simeq \pi X \quad \text{or} \quad t^N \pi X \simeq q^{-M} \pi X.$$

The second equation holds because the functors t and q commute with $t^N q^M$. If F has this form and $\mathcal{C}$ is monoidal then $\mathcal{C}/F$ is monoidal.

The proof of the main theorem found in section 4 shows that the modular group relations hold in a certain category $\mathcal{P}_{n,k}$ up to the application of a translation functor $\mathfrak{s}_k$. The equation in example 2.5 above will imply that the modular group acts on the associated orbit category $\mathcal{R}_{n,k} = \mathcal{P}_{n,k}/\mathfrak{s}_k$.

3. Braid groups and the modular group

In this section we recall the relationship between the braid group B_3 and the modular group $\mathrm{PSL}(2,\mathbb{Z})$. Notation established here will be used later.

3.1. **Braid groups.** If D_n is the closed disk with n punctures then the braid group B_n on n strands is the mapping class group

$$\mathrm{B}_n = \pi_0(\mathrm{Diff}^+(D_n, \partial)).$$

Alternatively, if $\mathrm{Config}_n(D^2)$ is the configuration space of n distinct points in the unit disk then the braid group B_n is the fundamental group,

$$\mathrm{B}_n = \pi_1(\mathrm{Config}_n(D^2), x_0),$$

see [1, 8]. Each loop $\gamma \in \mathrm{B}_n$ can be viewed as a braid in three space and each such braid can be represented by a diagram consisting of a sequence of crossings σ_i, $1 \leq i < n$. When $n = 3$, a presentation for the braid group in terms of these generators is given by

$$\mathrm{B}_3 = \langle \sigma_1, \sigma_2 : \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle.$$

3.2. **Modular groups.** The modular group $\mathrm{SL}(2,\mathbb{Z})$ is the group of 2×2 matrices with integer coefficients and determinant 1,

$$\mathrm{SL}(2,\mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a,b,c,d \in \mathbb{Z} \text{ and } ad - bc = 1 \right\}.$$

It is also the mapping class group of the torus T, see [8]. If $I \in \mathrm{SL}(2,\mathbb{Z})$ is the identity matrix then the center $\mathrm{Z}(\mathrm{SL}(2,\mathbb{Z})) \subset \mathrm{SL}(2,\mathbb{Z})$ is the subgroup $\{I, -I\}$ and the projective modular group $\mathrm{PSL}(2,\mathbb{Z})$ is defined by the quotient

$$\mathrm{PSL}(2,\mathbb{Z}) = \mathrm{SL}(2,\mathbb{Z})/\{I,-I\}.$$

Choosing representatives for $\{I,-I\}$ yields a $\mathbb{Z}/2$ action on the torus with the quotient, $S^2_4 = T/(\mathbb{Z}/2)$, the four punctured sphere. The group $\mathrm{PSL}(2,\mathbb{Z})$ is the mapping class group of S^2_4.

It can be shown that $\mathrm{PSL}(2,\mathbb{Z})$ is isomorphic to the free product $\mathbb{Z}/2 * \mathbb{Z}/3$, or equivalently,

$$\mathrm{PSL}(2,\mathbb{Z}) \cong \langle a, b : a^2 = 1, b^3 = 1 \rangle.$$

3.3. **Relating** B_3 **and** $\mathrm{PSL}(2,\mathbb{Z})$**.** Denote by q_n the braid

$$q_n = \sigma_{n-1}\sigma_{n-2}\cdots\sigma_1 \in \mathrm{B}_n\,.$$

For instance,

$q_3 =$  .

Multiplying q_n by itself n times produces the full twist, $T_n = q_n^n$. The equation,

$$\sigma_i T_n = T_n \sigma_i \quad \text{for} \quad i = 1, \ldots, n-1,$$

is proven by dragging σ_i through from the bottom of T_n to the top of T_n. It follows that T_n is contained in the center $\mathrm{Z}(\mathrm{B}_n) \subset \mathrm{B}_n$ of the braid group. In fact, $\langle T_n \rangle = \mathrm{Z}(\mathrm{B}_n)$, see [5]. When $n = 3$, the following proposition tells us that the quotient $\mathrm{B}_n / \langle T_n \rangle$ is related to the modular group.

Proposition 3.4. *The quotient of* B_3 *by the subgroup generated by the full twist* T_3 *is isomorphic to the projective modular group* $\mathrm{PSL}(2,\mathbb{Z})$*.*

$$\mathrm{B}_3 / \langle T_3 \rangle \cong \mathrm{PSL}(2,\mathbb{Z})$$

Proof. If we set $a = \sigma_1\sigma_2\sigma_1$ and $b = \sigma_1\sigma_2$ then

$$\mathrm{B}_3 / \langle T_3 \rangle = \langle \sigma_1, \sigma_2 : \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2, T_3 = 1 \rangle \cong \langle a, b : a^2 = 1, b^3 = 1 \rangle \cong \mathrm{PSL}(2,\mathbb{Z}).$$

□

The groups $\mathrm{B}_n / \langle T_n \rangle$ are always related to mapping class groups of punctured spheres, see section 4.2 of [1].

4. The Construction

In this section we construct sequences of dg categories,

$$\mathcal{R}_{n,k} \quad \text{where} \quad n > 0,\ k = n, n-2, n-4, \ldots, n \,(\mathrm{mod}\, 2),$$

and prove that each category supports an action of the modular group. We begin by defining categories $\mathcal{P}_{n,k}$ as extensions of $\mathrm{Kom}^*(3n)$ from section 2.1. The categories $\mathcal{R}_{n,k}$ are obtained from $\mathcal{P}_{n,k}$ using the orbit category construction of section 2.5. In theorem 4.4 below it is shown that there exists an action of the group $\mathrm{PSL}(2,\mathbb{Z})$ on each category $\mathcal{R}_{n,k}$. This induces an action of the modular group on $\mathcal{R}_{n,k}$.

The first step is to define a chain complex $E_{n,k}$ consisting of a composition of projectors.

Definition 4.1. ($E_{n,k}$) Suppose that $P_{n,k}$ is the higher order projector of [7], let

$$E_{n,k} = (P_{n,k} \sqcup P_{n,k} \sqcup P_{n,k}) \otimes P_{3n,k}.$$

In terms of pictures,



where the top row consists of three parallel $P_{n,k}$ projectors and the bottom row is a single $P_{3n,k}$.

Definition 4.2. ($\mathcal{P}_{n,k}$) The category $\mathcal{P}_{n,k} \subset \mathrm{Kom}^*(3n)$ is the image of the functor $- \otimes E_{n,k}$.

Alternatively, $\mathcal{P}_{n,k}$ is the full dg subcategory of $\mathrm{Kom}^*(3n)$ consisting to objects which are homotopy equivalent to those of the form $C \otimes E_{n,k}$.

Each object in the category $\mathcal{P}_{n,k}$ is bigraded. One can shift either the t-degree of the chain complex or q-degree of each object using the functors $t^{\frac{1}{2}}$ or q. There is a shift functor

$$\mathfrak{s}_k : \mathcal{P}_{n,k} \to \mathcal{P}_{n,k} \quad \text{where} \quad \mathfrak{s}_k = t^{k^2} q^{2k}.$$

In particular, if $C \in \mathcal{P}_{n,k}$ is a chain complex then the shift functor $\mathfrak{s}_k$ satisfies

$$\deg_t(\mathfrak{s}_k(C)) = \deg_t(C) + k^2 \quad \text{and} \quad \deg_q(\mathfrak{s}_k(C)) = \deg_q(C) + 2k.$$

These numbers are determined by property 2 in section 2.2.

The categories $\mathcal{R}_{n,k}$ are obtained from the categories $\mathcal{P}_{n,k}$ by requiring the shift functor $\mathfrak{s}_k$ to act by identity.

Definition 4.3. ($\mathcal{R}_{n,k}$) The differential graded category $\mathcal{R}_{n,k}$ is the orbit category of the shift functor $\mathfrak{s}_k$ acting on $\mathcal{P}_{n,k}$.

$$\mathcal{R}_{n,k} = \mathcal{P}_{n,k}/\mathfrak{s}_k$$

Theorem 4.4. *The group* $\mathrm{PSL}(2,\mathbb{Z})$ *acts on the category* $\mathcal{R}_{n,k}$.

Proof. Using the presentation $\mathrm{PSL}(2,\mathbb{Z}) = \langle a, b : a^2 = 1, b^3 = 1\rangle$ of section 3.2, we will construct a homotopy homomorphism

$$\mathrm{PSL}(2,\mathbb{Z}) \to \mathrm{End}(\mathcal{R}_{n,k}) \quad \text{where} \quad g \mapsto F_g.$$

Associated to the generators a and b are the chain complexes determined by the tangles

$a =$  and $b =$  ,

where each line represents an n-cabled strand. There are functors, $F_a, F_b : \mathcal{R}_{n,k} \to \mathcal{R}_{n,k}$, given by tensoring on the left. To the identity element $1 \in \mathrm{PSL}(2,\mathbb{Z})$, we associate the functor F_1 where 1 is the diagram consisting of $3n$ vertical strands. For inverses, a^{-1} and b^{-1}, we use the functors F_a and $F_{b\otimes b}$. To a word $w \cdot w' \in \langle a, b\rangle$, we associate the functor $F_w \circ F'_w$. In this manner, we obtain a functor $F_g : \mathcal{R}_{n,k} \to \mathcal{R}_{n,k}$ for each word $g \in \langle a, b\rangle$.

We proceed by checking that the relations $a^2 = 1$ and $b^3 = 1$ hold. Since the braids a^2 and b^3 are both full twists with some framing dependency, it suffices to check that this full twist is homotopy equivalent to the identity object in $\mathcal{R}_{n,k}$.

Consider what happens when the tangles a^2 or b^3 are glued onto $E_{n,k}$. As framed tangles, the full twist T_3 is isotopic to the tangle pictured on the right below.



Using the relations of section 2, we can move the small $P_{n,k}$ projectors to the top and absorb the top twists.



$$\simeq \left(t^{\frac{1}{2}k^2}q^k\right)^3$$



Using the bottom $P_{3n,k}$ to absorb the middle twist leaves us with

$$\left(t^{\frac{1}{2}k^2}q^k\right)^3\left(t^{\frac{1}{2}k^2}q^k\right)^{-1}$$



$$= \mathfrak{s}_k$$



$$\simeq$$



.

The last equality follows because of the cyclic grading in $\mathcal{R}_{n,k}$. We have shown that the relations $a^2 = 1$ and $b^3 = 1$ hold up to homotopy. ☐

Several variations in the choices made above also yield $\mathrm{PSL}(2,\mathbb{Z})$ representations. For instance, we could have instead chosen to associate to a^{-1} and b^{-1} the functors $F_{a^{-1}}$ and $F_{b^{-1}}$. In the same spirit, one could act using under-crossings instead of over-crossings.

Remark. A special case of the theorem applies to the subcategories $\mathcal{K}_n = \mathrm{Kom}^*(n,n,n,n)$ of $\mathrm{Kom}^*(4n)$ (see section 6 of [6]). Objects in $\mathcal{K}_n$ are homotopy equivalent to convolutions of chain complexes of the form,

(4.1)



where D is a disjoint union of intervals interfacing between four projectors. Reducing the grading so that $\mathfrak{s}_n$ acts by identity gives a new category $\mathcal{K}_n/\mathfrak{s}_n$ and, after fixing one projector, we can act on the other three by n-cabled strands as in the proof above.

The category $\mathcal{K}_n$ is analogous to the vector space assigned to the four punctured sphere S^2_4 by SU(2) Chern-Simons theory. The quotient, $T \to S^2_4$, described in section 3.2, allows one to identify the SU(2) skein space for S^2_4 with the SO(3) skein space for the torus at level $r = 2(n+1)$.

Spin networks of the form (4.1) correspond to $\mathrm{U}_q\,\mathfrak{sl}(2)$-equivariant maps from the nth irreducible representation V_n to $V_n^{\otimes 3}$. The dimension of this space is $n+1$.

$$\dim \mathrm{Hom}_{\mathrm{U}_q\,\mathfrak{sl}(2)}(V_n, V_n^{\otimes 3}) = n+1$$

Theorem 6.1 of [6] implies that $\dim \mathrm{K}_0(\mathcal{K}_n) = n+1$ and so we view the categories $\mathcal{K}_n/\mathfrak{s}_n$ as a sequence of representations of increasing complexity. Similar observations hold for the other categories mentioned above.

The corollary below notes that the above theorem defines representations of the modular group $\mathrm{SL}(2,\mathbb{Z})$ via pullback.

Corollary 4.5. *The group* $\mathrm{SL}(2,\mathbb{Z})$ *acts on the category* $\mathcal{R}_{n,k}$.

Proof. To each $x \in \mathrm{SL}(2,\mathbb{Z})$ associate the functor $F_{\pi(x)}$ defined in the proof of the theorem where $\pi : \mathrm{SL}(2,\mathbb{Z}) \to \mathrm{PSL}(2,\mathbb{Z})$ is the homomorphism. □

The decomposition of $\mathrm{Kom}^*(n)$ with respect to the central object T_n is an ingredient above. Each higher order projector $P_{n,k}$ corresponds to a different eigenvalue, or "frequency," of the full twist T_n (see property 2 section 2.2). In [7], each object $X \in \mathrm{Kom}^*(n)$ is shown to admit a decomposition in terms of these projectors. This implies that all objects are convolutions of complexes which respect the eigenvalue decomposition of the full twist. The Postnikov decomposition acts as a kind of Fourier transform. Passing to an orbit category removes some frequencies from consideration by introducing a periodicity.

5. A Concrete Example

In this section we consider the case when $n = 1$ and $k = 1$. We compare our categorical representation to a $\mathbb{C}$-vector space representation. The shorthand, $E = E_{1,1}$, $\mathcal{P} = \mathcal{P}_{1,1}$, $\mathfrak{s} = \mathfrak{s}_1$ and $\mathcal{R} = \mathcal{R}_{1,1}$, will be used.

We begin by considering the category $\mathrm{Kom}^*(3)$. The objects of $\mathrm{Kom}^*(3)$ are chain complexes of objects



which we denote by $1_3, e_1, e_2, e_1e_2$ and e_2e_1 respectively.

According to definition 4.2 in section 4, the category $\mathcal{P}$ consists of objects of the form $X \otimes E$ where $X \in \mathrm{Kom}^*(3)$ and E is the higher order projector $P_{3,1}$. From [6, 7], the projector $P_{3,1}$ is the chain complex $(P_{3,1}, d)$ where

$$P_{3,1} = \bigoplus_{n>0} t^{2n-1} q^{3n-2}(e_1 \oplus e_2) \oplus t^{2n} q^{3n-1}(e_1e_2 \oplus e_2e_1),$$

and the differential is given by

$$d = \begin{pmatrix} E & B \\ D & C \end{pmatrix}.$$

The maps in d are determined up to sign by the requirements in theorem 2.3 section 2.2 and are given explicitly in section 4.2 of [6].

For each $X \in \text{Pre-Cob}(3)$, thought of as an object in $\mathrm{Kom}^*(3)$, we have

$$X \otimes P_{3,1} = \begin{cases} X & \text{if } X \neq 1_3 \\ P_{3,1} & \text{if } X = 1_3 \end{cases}$$

so that the category $\mathcal{P}$ can be thought of as the subcategory consisting of chain complexes which do not include the identity object 1_3.

In order to define the representation of $\mathrm{PSL}(2,\mathbb{Z}) = \mathrm{B}_3 / \langle T_3 \rangle$, the chain complexes,

$S_1 =$  and $S_2 =$  ,

are associated to the generators $\sigma_1, \sigma_2 \in \mathrm{B}_3$. Tensoring on the left, $X \mapsto S_i \otimes X$, allows us to define two functors $F_1, F_2 : \mathcal{P} \to \mathcal{P}$. By setting $F_a = F_1 \circ F_2 \circ F_1$ and $F_b = F_1 \circ F_2$, the proof of theorem 4.4 section 4 shows that the functors F satisfy the relations necessary for a $\mathrm{PSL}(2,\mathbb{Z})$ action up to an application of $\mathfrak{s}$.

By passing to the orbit of $\mathfrak{s}$, we obtain the category $\mathcal{R}$ which satisfies, $\mathrm{Ob}(\mathcal{R}) = \mathrm{Ob}(\mathcal{P})$ and

$$\mathrm{Hom}_{\mathcal{R}}(X, Y) = \bigoplus_{m \geq 0} \mathfrak{s}^m \, \mathrm{Hom}^*(X, Y).$$

If $\pi : \mathcal{P} \to \mathcal{R}$ is the canonical projection then $\pi \circ \mathfrak{s} \simeq \pi$ implies that $\mathcal{R}$ supports an action of $\mathrm{PSL}(2,\mathbb{Z})$.

The Grothendieck group $\mathrm{K}_0(\mathcal{P})$ is spanned by the four objects e_1, e_2, e_1e_2 and e_2e_1. Using the convention that $\mathrm{K}_0(q) = -q$, $\mathrm{K}_0(t^{\frac{1}{2}}) = q^{\frac{1}{2}}$, the functors F_1 and F_2 yield matrices

$$[F_1] = \begin{pmatrix} -q^{-\frac{3}{2}} & 0 & q^{-\frac{1}{2}} & 0 \\ 0 & q^{\frac{1}{2}} & 0 & 0 \\ 0 & 0 & q^{\frac{1}{2}} & 0 \\ 0 & q^{-\frac{1}{2}} & 0 & -q^{-\frac{3}{2}} \end{pmatrix} \quad \text{and} \quad [F_2] = \begin{pmatrix} q^{\frac{1}{2}} & 0 & 0 & 0 \\ 0 & -q^{\frac{3}{2}} & 0 & q^{-\frac{1}{2}} \\ q^{-\frac{1}{2}} & 0 & -q^{\frac{3}{2}} & 0 \\ 0 & 0 & 0 & q^{\frac{1}{2}} \end{pmatrix}$$

In order to satisfy the relation $\pi \circ \mathfrak{s} \simeq \pi$, we proceed by setting $q = e^{\frac{2\pi i}{3}}$ and, in doing so, obtain a vector space

$$R = \mathrm{K}_0(\mathcal{P}) \otimes_{\mathbb{Z}[[q^{\frac{1}{2}}]][q^{-\frac{1}{2}}]} \mathbb{C}$$

comparable to the representation $\mathcal{R}$. The matrices $a = [F_1][F_2][F_1]$ and $b = [F_1][F_2]$ are given by,

$$a = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \quad \text{and} \quad b = \begin{pmatrix} 0 & 0 & -q & 0 \\ 0 & -q^{-1} & 0 & 1 \\ 1 & 0 & -q^{-1} & 0 \\ 0 & -q & 0 & 0 \end{pmatrix}.$$

These matrices define a $\mathrm{PSL}(2,\mathbb{Z})$ action on R. A computation shows that the relations $[F_1]^6 = 1$ and $[F_2]^6 = 1$ are also satisfied and so our representation factors through a smaller group,

$$G = \langle \sigma_1, \sigma_2 : \sigma_1^6 = \sigma_2^6 = 1, \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle.$$

So R is not a faithful representation of $\mathrm{PSL}(2,\mathbb{Z})$. On the other hand, the chain complexes $P_{3,1}$ and $F^6_{\sigma_i} \otimes P_{3,1}$ are unlikely to be homotopy equivalent.

References

[1] J. Birman, *Braids, links and mapping class groups*, Ann. Math Studies 82, Princeton University Press, 1975.

[2] A. I. Bondal and M. M. Kapranov, *Enhanced Triangulated Categories*, Math. USSR. Sbornik, vol. 70, 1991, no. 1, 93–107.

[3] D. Bar-Natan, *Fast Khovanov Homology Computations*, Journal of Knot Theory and Its Ramifications, vol. 16, 2007, no. 3, 243–255.

[4] D. Bar-Natan, *Khovanov's homology for tangles and cobordisms*, Geom. Topol., vol. 9, 2005, 1443–1499.

[5] W.-L. Chow, *On the algebraic braid group*, Ann. of Math., vol. 49, 1948, 654–658.

[6] B. Cooper and V. Krushkal, *Categorification of the Jones-Wenzl Projectors*, Quantum Topology, vol. 3, 2012, no. 2, 139–180.

[7] B. Cooper and M. Hogancamp, *An Exceptional Collection For Khovanov Homology*, arXiv:1209.1002.

[8] B. Farb and D. Margalit, *A Primer on Mapping Class Groups*, Princeton University Press, 2011.

[9] I. Frenkel, C. Stroppel and J. Sussan, *Categorifying fractional Euler characteristics, Jones-Wenzl projector and $3j$-symbols with applications to Exts of Harish-Chandra bimodules*, Quantum Topology, vol. 3, 2012, no. 2, 181–253.

[10] B. Keller, *On Triangulated Orbit Categories*, Documenta Math., vol. 10, 2005, 551–581.

[11] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J., vol. 101, 2000, no. 3, 359–426.

[12] M. Khovanov, *A functor-valued invariant of tangles*, Algebr. Geom. Topol., vol. 2, 2002, 665–741.

[13] M. Khovanov and P. Seidel, *Quivers, Floer cohomology, and braid group actions* J. Amer. Math. Soc., vol. 15, 2002, no. 1, 203–271.

[14] R. Lipshitz, P. Ozváth and D. Thurston, *A faithful linear-categorical action of the mapping class group of a surface with boundary*, Journal of European Mathematical Society, to appear.

[15] L. Rozansky, *An infinite torus braid yields a categorified Jones-Wenzl projector*, arXiv:1005.3266.

[16] L. Rozansky, *Khovanov homology of a unicolored B-adequate link has a tail*, arXiv:1203.5741.

[17] J. Roberts, *Rozansky-Witten Theory*, Proceedings of Georgia Topology conference, 2001, arXiv:math.QA/0112209.

[18] R. Rouquier, *Categorification of sl2 and braid groups*, Trendsin representation theory and related topics, Amer. Math. Soc., 2006, 137–167.

[19] P. Seidel and R. Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J., vol. 108, 2001, no. 1, 37–108.

Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich

E-mail address: `benjamin.cooper@math.uzh.ch`